

## Ultrastructural aspects of *Eriogaster catax* and *Eriogaster lanestris* (Lepidoptera: Lasiocampidae)

Cristian Sitar<sup>1</sup>✉ and Alexandra Ciorîță<sup>1, 2</sup>

<sup>1</sup>Babeș-Bolyai University, Faculty of Biology and Geology, 5-7 Clinicilor, 400006, Cluj-Napoca, Romania; <sup>2</sup>National Institute for Research and Development of Isotopic and Molecular Technologies, 67-103 Donat, 400293 Cluj-Napoca, Romania;  
✉Corresponding author, E-mail: cristiansitar@yahoo.com.

**Abstract.** *Eriogaster catax* (Linnaeus, 1758) and *E. lanestris* (Linnaeus, 1758) are two related species in the family Lasiocampidae with decreasing distribution areas and extremely isolated populations, thus being vulnerable to numerous threats, particularly human impact. Until presently, the biological and ecological studies regarding the species in question are scarce, particularly for *E. catax*, a species protected through Annexes II and IV of the Council Directive 92/43/EEC, Annex II of the Bern Convention and Law no. 49/2011 in Romania. Taking into account the importance of the *E. catax* species and the insufficient information reported on the ecological need of the species, a highly detailed analysis at an ultrastructural level could provide valuable information. The primitive aspect of the wings explained the weak distribution of adults. The different flight period of adults and egg laying season were reflected on the structure, adherence and size of eggs. Also, the structure of the fibers was correlated with the different ways in which *E. catax* and *E. lanestris* use the nest.

**Keywords:** *Eriogaster catax*, *E. lanestris*, oviposition strategy, ecological niche.

### Introduction

*Eriogaster catax* (Linnaeus, 1758) and *E. lanestris* (Linnaeus, 1758), (Lepidoptera: Lasiocampidae) are two cohabitant species (Sáfián, 2006) which are mainly found in the semi-natural habitats that have a predominant shrubby vegetation (Kadej *et al.*, 2018).

*E.catax* is one of the least studied species due to living in very localized areas (Drews and Wachlin, 2003) and it is also considered an endangered species in several European countries. Currently it is protected by law at European level by the *Directive 92/43/EEC*, Annexes II and IV and the *Berne Convention*, Annex II

(Höttinger, 2005), while also being categorized as data deficient (DD) in the IUCN red list of threatened species (Farkač *et al.*, 2005). The species spreads throughout entire Europe, starting from the north of Spain up to the Balkans, and to the south down to the Ural Mountains and Anatolian peninsula (Baillet, 2013; Borges, 2012; Bury, 2015; Freina, 1996; Freina and Witt, 1987; Karsholt and Razowski, 1996; Konvička *et al.*, 2005; Leraut, 2006; Ruf *et al.*, 2003).

*E. lanestris* has a much larger distribution and can be found in almost all Europe except for the Tundra habitats, the Mediterranean region and in Asia with reports up to north of Caucasus, Kazakhstan, Siberian southern region, Central Yakutia and the Amur basin (Ruf *et al.*, 2003; Ebert, 1994; Freina and Witt, 1987; Dubatolov and Zolotuhin, 1992; Pro Natura, 2005).

In their larval stage, both species feed primarily on the *Prunus spinosa* and *Crataegus monogyna* shrubs. The female moths of both species lay their eggs only once, on the branches of the host plant (Baillet, 2011; Caron, 2009; Höttinger, 2005; Oleksa, 2010). Due to the absence of a "mouth", the adult moths do not feed in their life span and rely solely on their larval nutrition (Malkiewicz, 2015).

A distinguishing feature of the two species is their highly social behavior during the larval stage, categorized as gregarious (Baillet, 2013; Bury, 2015; Caron, 2009; Chrzanowski *et al.*, 2013; Freina, 1996; Ruf *et al.*, 2003).

In the literature published so far, the growth stages of *E. catax* and *E. lanestris* have been described from a morphological point of view. By using a scanning electron microscopy technique, Fitzgerald (1995) highlights a series of ultrastructural characteristics such as the texture of the silk nest woven by the caterpillars or different types of glands from the genus *Malacosoma*, which is related to the genus *Eriogaster*.

Taking into account the importance of the *E. catax* species and the scarce information reported on the ecological need of the species, a highly detailed morphological analysis at an ultrastructural level could provide valuable information. Due to the fact that *E. catax* and *E. lanestris* share the same habitat and sometimes even the same shrub during their larval development stage, the investigation of both species was considered necessary.

## **Materials and methods**

### ***Electron microscopy analysis***

Samples (adults, eggs, larva and nest silk) were prepared for scanning electron microscopy (SEM) analysis using the turbomolecular pumped coater Quorum Q150T ES, from the Integrated Electron Microscopy Laboratory (LIME) of National Institute for Research and Development of Isotopic and Molecular Technologies (INCDTIM), Cluj-Napoca, Romania, and examined at SEM Hitachi SU8230 (LIME-INCDTIM, Cluj-Napoca, Romania).

### **Statistics**


Statistical analyses were conducted using online *t* test calculator.

### **Results and discussions**

#### **Adults**

Both species had palmate scales (Fig. 1 A, B, D), fixed on the surface of the wings through a peduncle (Fig. 1 C), and were randomly dispersed on the surface.

The fact that the scales were fixed through a peduncle indicates that they originate from trichomes. Some studies showed the relation between Lepidoptera scales and sensory trichomes (Galant *et al.*, 1998; Zhou *et al.*, 2009). The scales distribution of the scales is correlated with the underdeveloped flight ability, explaining also the low dissemination of females. Peduncle-like scales and no mouthpieces are primitive characters of *Eriogaster*.



**Figure 1.** A. Palmate scales of *E. catax*; B. Palmate scales of *E. lanestris*; C. The insertion point of a peduncle to the scale; D. The surface of a scale in detail.

The antennae of both sexes were consisted of three segments: scape, pedicel and a long (Ma *et al.*, 2017; Mark *et al.*, 2017) bipectinate flagellum (Fig. 2 A). The bipectinate flagellum is composed of approximately same number of flagellomeres in males and females. Each flagellomere bears two lateral branches (Fig. 2 C, 4 A). The females of both species had shorter and thicker branches than the males. There are also differences in shape and thickness between the flagellomere branches of the two species i.e., those of *E. lanestris* were significantly thickened towards the base giving them a pear-like shape (Fig. 2 C, D).

Branches of flagellomeres in males of the two species increased progressively in length from the proximal end to the middle, and then decreased towards the apex. Branch length was sexually dimorphic, with males possessing significantly longer branches throughout.

The dorsal surface of the flagellum (antennal spindle) and the entire scape and pedicel were covered in overlapping lamellar scales (Fig. 2 B) (Mark *et al.*, 2017; Yuvaraj *et al.*, 2018).

Based on the external morphology of the sensilla observed under SEM, many sensilla with role in olfaction which plays a critical role for insects were identified (Yuvaraj *et al.*, 2016).

Sensilla trichodea were predominantly distributed on the ventral side of the lateral branches (Fig. 3A, B; 4A) (Fernandes *et al.*, 2017; Li *et al.*, 2018; Mark *et al.*, 2017). In males they were grouped in clusters of 5–6 sensilla (Fig. 4 B, C, D). Sensilla trichodea are the most abundant sensory receptors on the antennae of both sexes and species (Fig. 3 A, B and 4 A).

On the dorsal side of the lateral branches (Fig. 2 C, D; 4 A) we identified 8-9 (n=100) sensilla chaetica (Fig. 3 E, F and 4 E, F) (Mark *et al.*, 2017) at antennae of both species.


On the apex of of the lateral branches of both sexes we identified 1-3 sensilla basiconica (Fig. 3 C, D) (Mark *et al.*, 2017).

### **Eggs**


For *E. catax* the eggs had an elongated, cylindrical form, nicely organized in rows and very well welded together and to the substrate. On the apex, the eggs had the micropylar region formed like a small dent, with numerous pores (Fig. 5 A, C, E).

*E. lanestris* had less organized eggs, with weak connections in between and to the surface. The eggs were elongated and with a wider apex compared with the basal part. The micropylar region is slightly curved (Fig. 5 B, D, F). Because the eggs are not so close to each other, the pores developed on the lateral of the egg, and not on top, like *E. catax* (Fig. 6 A).


Both species hatch by penetrating the micropylar region and the missing part indicated that the caterpillars consumed this region which contributed to their nutrition (Fig. 6 B).




**Figure 2.** SEM micrographs showing the external morphology of the antennae of *E. catax* and *E. lanestris*: A. The bipectinate antennae of the *E. catax* female; B. The dorsal surface of the flagellum is covered with overlapping lamellar scales at both species; C. Female antenna of *E. catax* - lateral view. D. Female antenna of *E. lanestris* - lateral view; E. Female antenna of *E. catax* - ventral view. F. Female antenna of *E. lanestris* - ventral view;




**Figure 3.** The external morphology of the sensilla observed using SEM on the female antennae of *E. catax* (A., C., E.) and *E. lanestris* (B., D., F.). A., B. Sensilla trichodea on the ventral side of the lateral branches; C., D. Sensilla basiconica on the apex of the lateral branches; E., F. Sensilla chaetica on the dorsal side of the lateral branches.



**Figure 4.** The external morphology of the antennae and the sensilla observed on SEM on the male's antennae of *E. catax*. A. Male antenna of *E. catax* - ventral view. Arrows indicate the location of the chaetica sensilla. B., C. Sensilla trichodea on the ventral side of the lateral branches of both species; They were grouped in clusters of 5–6 sensilla; E. Sensilla chaetica on the dorsal side of the lateral branches. F. Sensilla basiconica on the apex of of the lateral branches.



**Figure 5.** A. Eggs of *E. catax* arranged in ordered rows. The eggs are tightly bound together and adhere very well to the substrate; B. Eggs of *Eriogaster lanestris* - the rows are more slightly ordered and the adhesion to the substrate is weak; C. Around the micropillary region of the *E. catax* egg, numerous pores indicated by arrows can be observed; D. The micropillary region of the *E. lanestris* egg is slightly bulged like a dome. E. Lateral view of *E. catax* eggs; F. Lateral view of *E. lanestris* eggs. Arrows indicate the contact area with adjacent eggs.



**Figure 6.** A. Respiratory pores on the egg surface of *E. lanestris*;  
B. Hatch orifices of *E. lanestris* caterpillars.

*E. lanestris* has smaller eggs compared to *E. catax* and this could be explained by the different time periods when the eggs are laid. *E. catax* lays its eggs in September-October (Bury, 2015; Chrzanowski et al., 2013; Freina, 1996), and *E. lanestris*, during March-April (Ruf et al., 2003).

Compact and well adhered eggs on the surface of the leaves, along with a small number of large eggs are adaptations of *E. catax* to overcome unfavorable conditions during winter period.

### **Larva**

The polypod caterpillar of *E. catax* had five stemmata (the visual organs of the larva) on each side of the cephalic capsule. Also, numerous pores were observed on the cephalic capsule most probably act like atmospheric pressure sensors (Fig. 7 A, B, C).


Both species had protective tubular trichomes on the surface of the body and on the cephalic capsule. To evaluate the differences, trichomes' width on the cephalic capsule were measured and very significant differences were observed ( $p < 0.05$ ). *E. catax* had an average width of trichomes of  $8.96 \mu\text{m} \pm 0.2 \mu\text{m}$  ( $N=30, \pm \text{SEM}$ ) and for *E. lanestris* the trichomes measured  $4.51 \mu\text{m} \pm 0.25 \mu\text{m}$  ( $N=30, \pm \text{SEM}$ ) (Fig. 4 D, E, F).

On the sides of the body, eight pairs of spiracles were observed. Spiracles contain numerous atrial structures that close or open the airway. The atrial structures of the spiracles are finger-shaped and they have many cilia (Fig. 8. A, B).


On the ventral side the larva had three pairs of thoracic legs and four pairs of prolegs provided with numerous crochets (Fig. 9).

*E. catax* hatch in early spring, along with bud break of *Prunus spp.* and *Crataegus spp.* (Baillet, 2013; Caron, 2009; Höttinger, 2005).


Access to food is still limited during this period and cold weather in the morning and at night lead to high consumption of energy for thermoregulation. *E. lanestris* hatches 3 to 4 weeks later (Konvička *et al.*, 2006), when both *Prunus* spp. and *Crataegus* spp. are blooming, the temperatures slightly increased and food sources are plenty.



**Figure 7.** A. Cephalic capsule of the caterpillar; B. Five stemmata - the visual organs of the larva; C. Cephalic capsule of the caterpillar has many pores - atmospheric pressure sensors; D. Protective trichomes present on the body and on the cephalic capsule; E. Protective trichome - cross section; F. *E. catax* average width of trichomes - 8.96  $\mu\text{m}$ .




**Figure 8.** Two different types of caterpillars legs. A., B. Three pairs of articulated abdominal legs C., D. Proleg provided with numerous crochets.



**Figure 9.** Spiracle with atrial structure. A. Caterpillar in first instar *L1*;  
B. Caterpillar in last instar *L5*.

### ***Nest silk***

*E. catax* had curvy-like fibers, welded together in different spots, and *E. lanestris* had straight fibers which did not form agglomerations, and with calcium oxalate crystals for increased strength. The width of the fibers was measured and highly significant differences were observed ( $p < 0.001$ ). *E. catax* had  $0.89 \mu\text{m} \pm 0.04 \mu\text{m}$  thick fibers, and *E. lanestris* had  $1.78 \mu\text{m} \pm 0.07 \mu\text{m}$  ( $N = 100, \pm\text{SEM}$ ), making it double in width (Fig. 10).



**Figure 10.** The ultrastructural aspects of *E. catax* (A, C, E) and *E. lanestris* (B, D, F) silk

The small width of the fibers of *E. catax* is compensated by their capacity to weld and form a felt-like structure, which confers rigidity to the nest. For *E. lanestris* the time required for the fibers to stiffen is shorter compared to *E. catax*. This explains why for *E. catax* the fibers adhere to each other.

The nest made by *E. catax* is used as a platform to rest on, gain warmth and shelter. The nest grows until the larva reach L3 stage, but they do not have enough space to stay inside it, compared to *E. lanestris*, where the larva can get inside the nest.

The nest serves as an activity center and plays an important role in the thermoregulation of the colony (Fitzgerald and Costa, 1999; Costa, 1997).

## Conclusions

SEM analysis revealed precious information about the ecology and biology of the species. The primitive aspect of the wings explained the weak spatial distribution of the adults. The different flight period of adults and egg laying season were reflected in the structure, adherence and size of eggs. Also, the structure of the fibers was influenced by the ways in which *E. catax* and *E. lanestris* use the nest.

**Acknowledgements.** We gratefully acknowledge to „Constantin Craciun” Electron Microscopy Laboratory and the Integrated Electron Microscopy Laboratory (LIME) of the National Institute for Research and Development of Isotopic and Molecular Technologies (INCDTIM), Cluj-Napoca, Romania.

## REFERENCES

- Baillet, Y. (2011). Inventaire de *Eriogaster catax* (Laineuse du Prunellier) sur l'ENS des communaux de Trept. *Rapport d'étude rendu au format Excel de Flavia, Trept.*
- Baillet, Y. (2013). Inventaire de *Eriogaster catax* (Laineuse du Prunellier) sur l'ENS des communaux de Trept. *Rapport d'étude rendu au format Excel de Flavia, Trept.*
- Borges, A. (2012). La Laineuse du prunellier – 2.localisation des chenilles (dans les nids communautaires) : c'est maintenant ! I NSECTES n°164 - 2012 (1)
- Bury J. (2015). Nowe dane o występowaniu *Eriogaster catax* (Linnaeus, 1758) (Lepidoptera: Lasiocampidae) w południowo-wschodniej Polsce. *Acta Ent. Siles.* 23 (online 012)1–10.

- Caron, G. (2009). La laineuse du prunellier *Eriogaster catax* (Linnaeus, 1758) (Lepidoptera, Lasiocampidae) victime des changements climatiques ? Ecologie de l'espèce et hypothèses sur son déclin dans la région genevoise. *Entomo Helvetica*, 2: 49-60.
- Chrzanowski, A., Kuźmiński, R., Łabędzki, A., Mazur, A., & Rutkowski, P. (2013). Occurrence of *Eriogaster catax* (Linnaeus, 1758) (Lasiocampidae, Lepidoptera) and the proposed protective actions on the Polish territory. *Nauka Przyroda Technologia* 7, 2, (24): 1-7.
- Costa, J.T. (1997). Caterpillars as social insects. *American Scientist*: 85, 150-159.
- Drews, M., & Wachlin, V. (2003). *Eriogaster catax* (Linnaeus, 1758). W: Petersen B., Ellwanger G., Biewald G., Hauke U., Ludwig G., Pretscher P., Schröder E., Ssymank A. (red.). Das europäische Schutzgebietssystem Natura 2000. Ökologie und Verbreitung von Arten der FFH-Richtlinie in Deutschland. Band 1: Pflanzen und Wirbellose. Schr.R. Landschaftspflege Naturschutz 69/1: 459.464.
- Dubatolov, V.V., & Zolotuhin, V.V. (1992). A list of the Lasiocampidae from the territory of the former USSR. *Atalanta* 23 (3/4): 531-548.
- Ebert, G. (1994). Die Schmetterlinge Baden-Württembergs. Band 4, Nachtfalter II. Ulmer Verlag, Stuttgart.
- Farkač, J., Král D., & Škorpík, J. (2005). Červený seznam ohrožených druhů České republiky Bezobratlí. Agentura ochrany přírody a krajiny ČR. 760pp
- Fernandes, F.F., & Pimenta, P.F.P. (2018). Typology and distribution of antennal sensory organs of adults *Haematobia irritans* of the Midwest Region of Brazil. *Pesquisa Veterinária Brasileira*, 38(1), 183-188. doi:10.1590/1678-5150-pvb-5499
- Fitzgerald, T.D. (1995). The tent caterpillars. Cornell University Press. ISBN 0-8014-2456-9
- Fitzgerald, T.D., & Costa J.T. (1999). Collective behavior in social caterpillars. pp. 379-400 in Detrain C, Deneubourg JL and Pasteels JM (eds.) Information processing in social insects. Birkhäuser Verlag, Basel.
- Freina, J.J. de, (1996). *Eriogaster catax* (Linnaeus, 1758). W: Helsdingen P.J., van Willemse L., Speight M.C.D. (red.): Background information on invertebrates of the Habitat Directive and the Bern Convention. Part I: Crustaceae, Coleoptera and Lepidoptera. *Nature and Environment* No. 79: 117.120.
- Freina, J.J. de, & Witt, T.J. (1987). Die Bombyces und Sphinges der Westpaläarktis (Insecta, Lepidoptera). Edition Forschung und Wissenschaft, München.
- Galant, R., Skeath, J.B., Paddock, S., Lewis, D.L., & Carroll, S.B. (1998). Expression pattern of a butterfly achaete-scute homolog reveals the homology of butterfly wing scales and insect sensory bristles. *Current Biology* 1998, 8:807-813.
- Höttinger, H. (2005). Der Hecken-Wollflügel (*Eriogaster catax* L.) in Wien (Lepidoptera: Lasiocampidae). Endbericht einer Studie im Auftrag der Wiener Magistratsabteilung MA 22 (Umweltschutz).
- Kadej, M., Zajac, K., & Tarnawski, D. (2018). Oviposition site selection of a threatened moth *Eriogaster catax* (Lepidoptera: Lasiocampidae) in agricultural landscape - implications for its conservation. *Journal of Insect Conservation*, 22, 29-39.

- Karsholt, O. & Razowski J. (eds.). (1996) The Lepidoptera of Europe. A distributional checklist. Apollo Books, Stenstrup.
- Konvička, M., Beneš, J., & Čížek, L. (2005). Ohrožený hmyz nelesních stanovišť: ochrana a management. *Sagittaria* 2005. 127 pp.
- Konvička, M., Hula, V., & Beneš, J. (2006). Metodika monitoringu evropsky významného druhu bourovec trnkový (*Eriogaster catax*). Nепublikovaný rukopis, deponován na UP AOPK ČR Praha. 7pp.
- Leraut, P. (2006). Moths of Europe. Volume 1: Saturnids, Lasiocampids, Hawkmoths, Tiger Moths. N.A.P. Editions, Verrières le Buisson. 395 pp.
- Li, Y., Liu, F., Du, X., Li, Z., & Wu, J. (2018). Ultrastructure of antennal sensilla of three fruit borers (Lepidoptera: Crambidae or Tortricidae). *PLoS ONE* 13(10): e0205604. doi:10.1371/journal.pone.0205604
- Ma, M., Chang, M., Lu, Y., Ley, C., & Yang, F. (2017). Ultrastructure of sensilla of antennae and ovipositor of *Sitotroga cerealella* (Lepidoptera: Gelechiidae), and location of female sex pheromone gland. *Scientific Reports* 7, 40637. doi:10.1038/srep40637
- Malkiewicz, A. (2015). Barczatka kataks w Krainie Łęgów Odrzańskich – występowanie, zagrożenia, zalecenia ochronne. In: Pietkiewicz M. and Sztwiertnia H. 2015 Kronika Łęgów Odrzańskich nr 1. Rzadkie i chronione zwierzęta krainy Łęgów Odrzańskich - zagrożenia i perspektywy rozwoju: Zimowanie ptaków w Krainie Łęgów Odrzańskich. Prochowice 2015. ISBN: 978-83-942200-0-6
- Mark, C., Parsons, S., & Holwell, G. (2017). Antennal morphology and micro-sensory architecture of the New Zealand magpie moth, *Nyctemera annulata* (Lepidoptera: Erebiidae): diversity, distribution and dimorphism. *Austral Entomology*. doi: 10.1111/aen.12279
- Oleksa, A. (2010). 1075 Barczatka kataks *Eriogaster catax* (Linnaeus, 1758) [Eastern eggar *Eriogaster catax* (Linnaeus, 1758)]. In: Monitoring gatunków i siedlisk przyrodniczych ze szczególnym uwzględnieniem specjalnych obszarów ochrony siedlisk Natura 2000. Wyniki monitoringu. GIOŚ Warszawa [in Polish].
- Pro Natura (2005). Lepidopterologen-Arbeitsgruppe (2000): Schmetterlinge und ihre Lebensräume, Band 3. Pro Natura - Schweizerischer Bund für Naturschutz, ISBN 3855870322, 928 Seiten.
- Ruf, C., & Fiedler, K. (2002). Plasticity in foraging patterns of colonies of the small eggar moth, *Eriogaster lanestris* (Lepidoptera: Lasiocampidae). *Oecologia* 131: 626–634.
- Ruf, C., Freese, A., & Fiedler, K. (2003). Larval sociality in three species of central-place foraging lappet moths (Lepidoptera: Lasiocampidae): a comparative survey. *Zoologischer Anzeiger - A Journal of Comparative Zoology* 242 (3): 209.222.
- Sáfián, S. (2006). The occurrences of *Eriogaster catax* (Linnaeus, 1758) and *Eriogaster lanestris* (Linnaeus, 1758) (Lepidoptera: Lasiocampidae) in the Körös Valley (South-Eastern Hungary) *Folia Entomologica Hungarica*, 67, 137-143.
- Yuvaraj, J.K., Andersson, M., Anderbrant, O., & Löfstedt, C. (2018). Diversity of olfactory structures: A comparative study of antennal sensilla in Trichoptera and Lepidoptera. *Micron*, vol. 111, pp. 9-18. doi:10.1016/j.micron.2018.05.006

Zhou, Q., Yu, L., Shen, X., Li, Y., Xu, W., Yi, Y., & Zhang, Z. (2009). Homology of Dipteran Bristles and Lepidopteran Scales: Requirement for the *Bombyx mori* achaete-scute Homologue ASH2. *Genetics* 183: 619–627. doi: 10.1534/genetics.109.102848